

DISCRETENESS AND RATIONALITY OF F -THRESHOLDS

MANUEL BLICKLE, MIRCEA MUSTĂŢĂ, AND KAREN E. SMITH

ABSTRACT. The F -thresholds are characteristic p analogs of the jumping coefficients for multiplier ideals in characteristic zero. We give an alternative description of the F -thresholds of an ideal in a regular and F -finite ring R , which enables us to settle two open questions posed in [MTW]. Namely, we show that if the ring is, in addition, essentially of finite type over a field, then the F -thresholds are rational and discrete.

1. INTRODUCTION

In recent years, multiplier ideals have played an important role in higher dimensional birational geometry. For a given ideal $\mathfrak{a}$ on a smooth variety X , and a real parameter $c > 0$, the multiplier ideal $\mathcal{J}(\mathfrak{a}^c)$ is defined via a log resolution of the pair $(X, \mathfrak{a})$. Recall that a log resolution is a proper birational map $\pi : X' \rightarrow X$, with X' smooth such that $\mathfrak{a}\mathcal{O}_{X'}$ defines a simple normal crossing divisor $A = \sum_{i=1}^r a_i E_i$. Then, by definition,

$$(1) \quad \mathcal{J}(\mathfrak{a}^c) := \pi_* \mathcal{O}_{X'}(K_{X'/X} - \lfloor cA \rfloor),$$

and this is an ideal of $\mathcal{O}_X$ that does not depend on the chosen log resolution. A jumping coefficient (also called a jumping number or a jumping exponent) of $\mathfrak{a}$ is a positive real number c such that $\mathcal{J}(\mathfrak{a}^c) \neq \mathcal{J}(\mathfrak{a}^{c-\varepsilon})$ for every $\varepsilon > 0$. These invariants have been introduced and studied in [ELSV]. It follows from the formula (1) that if c is a jumping coefficient, then $c \cdot a_i$ is an integer for some i . In particular, every jumping coefficient is a rational number, and the set of jumping coefficients of a given ideal is discrete.

Hara and Yoshida introduced in [HY] a positive characteristic analog of multiplier ideals, denoted by $\tau(\mathfrak{a}^c)$. This is a generalized test ideal for a tight closure theory with respect to the pair $(X, \mathfrak{a}^c)$. Similarly, one can define jumping numbers for these test ideals. These invariants were studied under the name of F -thresholds in [MTW], where it was shown that they satisfy many of the formal properties of the jumping coefficients in characteristic zero.

2000 *Mathematics Subject Classification.* Primary: 13A35; Secondary: 14B05.

Key words and phrases. F -thresholds, test ideals, jumping coefficients, jumping exponents.

The first author was partially supported by the *DFG Schwerpunkt Komplexe Geometrie*. The second and third authors were partially supported by the NSF under grants DMS 0500127 and DMS 0500823, respectively.

We emphasize that the test ideals are not determined by a log resolution of singularities, even in the cases where such a resolution is known to exist. Instead, the definition uses the Frobenius morphism and requires a priori infinitely many conditions to be checked. This lack of built-in finiteness makes the question of rationality and discreteness of the F -thresholds non-trivial, and in fact these properties were left open in [MTW].

In the present paper we settle these questions in the case of a regular ring R that is essentially of finite type over an F -finite field: more precisely, we show that for every ideal in such a ring, all F -thresholds are rational and they form a discrete set. We mention that in the case when $\mathfrak{a}$ is a principal ideal in $R = k[[x, y]]$, where k is a finite field, this statement has been proved recently by Hara and Monsky in [HM]. Our key result, which implies discreteness and rationality of F -thresholds, is a finiteness statement for test ideals in a polynomial ring. We show in Proposition 3.2 below that if $\mathfrak{a}$ is an ideal in $k[x_1, \dots, x_n]$ that is generated in degree $\leq d$, then the test ideal $\tau(\mathfrak{a}^c)$ is generated in degree $\leq \lfloor c \cdot d \rfloor$.

The structure of the paper is as follows. In Section 2 we recall the definition of test ideals and F -thresholds in a slightly different setup from that in [HY] and [MTW]. In fact, we prefer to work with a different definition of test ideals that is more suitable for our purpose, and show later that this is equivalent with the definition from [HY]. Due to this alternative definition, and to the slightly different setup, we decided to develop the theory from scratch. We hope that this would be beneficial for the reader, as with the present definition the basic results become particularly transparent. After this initial setup, we prove our main results in Section 3.

1.1. Acknowledgements. We are grateful to Rob Lazarsfeld for useful discussions, to Craig Huneke for a suggestion regarding the notation in the paper, and to the referee for several helpful comments.

2. GENERALIZED TEST IDEALS AND F -THRESHOLDS

In this section we collect some basic results about generalized test ideals as introduced by Hara and Yoshida [HY] (see also [Tak2] and [HT]). We restrict ourselves to working over an F -finite regular ring which allows us to use an alternative description of these ideals. We subsequently prove that our ideals agree with the ones defined in [HY]. An advantage of our definition is that the basic properties are very easy to prove. We include some of the characteristic p analogs of some of the basic results on multiplier ideals, such as the Restriction Theorem (Remark 2.28), Skoda's Theorem (Proposition 2.25) or the Subadditivity Theorem (Proposition 2.11(4)), all with very elementary proofs.

We also recall the definition of F -thresholds from [MTW], though we do not restrict to the case of a *local* ring as in *loc. cit.* In this more general framework we show that the set of F -thresholds coincides with the set of jumping exponents for the test ideals.

Let us fix the following notation: R denotes a regular F -finite ring of positive characteristic p . We stress here that R is not assumed to be local. The regularity of R implies that the Frobenius morphism $F: R \rightarrow R$ sending $x \in R$ to x^p is flat [Ku]. This is equivalent to saying that the functor $(F^e)^*$ of extending scalars via F^e is exact. By our assumption that R is F -finite we just mean that F is a finite morphism. Since F is also flat, it follows that R is a finitely generated locally free module over its subring R^p of p^{th} powers.

For an ideal J of R and a positive integer e we set $J^{[p^e]} := (u^{p^e} | u \in J)$. If $J = (u_1, \dots, u_r)$, then $J^{[p^e]} = (u_1^{p^e}, \dots, u_r^{p^e})$. Note that we have $(J^{[p^e]})^{[p^{e'}]} = J^{[p^{e+e'}]}$. One easily verifies that $(F^e)^*(R/J) \cong R/(J^{[p^e]})$. In particular, since F is (faithfully) flat we see that if u is in R , then $u^p \in J^{[p]}$ if and only if $u \in J$.

Example 2.1. For a field k , being F -finite precisely means that $[k: k^p] < \infty$. One easily verifies, by explicitly giving a basis of R over R^p , that for such fields the polynomial ring $k[x_1, \dots, x_n]$ and even the power series ring $k[[x_1, \dots, x_n]]$ in finitely many variables are F -finite.

More generally, if A is an F -finite ring, then every A -algebra that is essentially of finite type over A is again F -finite. To check this observe that if S is a multiplicative system in a ring R , and if $a_1, \dots, a_N$ generate R over R^p , then $\frac{a_1}{1}, \dots, \frac{a_N}{1}$ generate $S^{-1}R$ over $(S^{-1}R)^p$.

Conversely, if k is a field and if R is an F -finite k -algebra essentially of finite type over k , then $[k: k^p] < \infty$. Indeed, if $\mathfrak{m}$ is a maximal ideal of R and $K = R/\mathfrak{m}$, then $[K: K^p] < \infty$. Since $[K: k] < \infty$, we deduce that $[k: k^p] < \infty$.

If $(R, \mathfrak{m})$ is a regular local ring that is F -finite, then its completion $\widehat{R}$ is also F -finite (and regular). In fact, since R is local, R is free over R^p . If $a_1, \dots, a_N$ give a basis of R over R^p , we claim that these elements also give a basis of $\widehat{R}$ over $\widehat{R}^p$. Indeed, we have canonical isomorphisms

$$F^*(\widehat{R}) = F^*(\varprojlim_{\ell} R/\mathfrak{m}^{\ell}) \simeq \varprojlim_{\ell} F^*(R/\mathfrak{m}^{\ell}) \simeq \varprojlim_{\ell} R/(\mathfrak{m}^{\ell})^{[p]} = \widehat{R}$$

(note that since F is finite, F^* commutes with the above projective limit). Therefore the Frobenius morphism on $\widehat{R}$ is obtained by base extension from the Frobenius morphism on R , which implies our claim.

2.1. The ideals $\mathfrak{b}^{[1/q]}$. Let $\mathfrak{b}$ be an ideal of R and let $q = p^e$. We now introduce certain ideals $\mathfrak{b}^{[1/q]}$ which are crucial for our definition of the generalized test ideals. More precisely, the test ideals of $\mathfrak{a}$ are constructed using all $(\mathfrak{a}^r)^{[1/p^e]}$. The ideals $(\mathfrak{a}^r)^{[1/p^e]}$ were introduced in the case of a principal ideal $\mathfrak{a}$ in [AMBL] to study the D -module structure

of a localization of R . It was predicted in [AMBL] that these ideals might be related to tight closure theory and the present work confirms this prediction. Furthermore, similar ideas also appear in the description of generalized test ideals in [HT, Lemma 2.1].

Definition 2.2. For an ideal $\mathfrak{b}$ of R and $q = p^e$, where e is a positive integer, let $\mathfrak{b}^{[1/q]}$ denote the unique smallest ideal J of R with respect to inclusion, such that

$$\mathfrak{b} \subseteq J^{[q]}.$$

Since R is a finitely generated flat (hence projective) R -module via F^e , the well-known lemma below implies that for every family of ideals $\{J_i\}_i$ of R we have $(\cap_i J_i)^{[q]} = \cap_i J_i^{[q]}$. Therefore $\mathfrak{b}^{[1/q]}$ is well-defined. We make the convention that $\mathfrak{b}^{[1/p^0]} = \mathfrak{b}$. Note in particular that $\mathfrak{b}^{[q'/q]}$ makes sense for every $q = p^e$ and $q' = p^{e'}$.

Lemma 2.3. *If M is a finitely generated projective module over the Noetherian ring R , then for every family $\{J_i\}_i$ of ideals of R we have*

$$\cap_i J_i M = (\cap_i J_i) M.$$

Proof. The assertion is clear when M is free. The general case follows since we can find a finitely generated free module P such that M is a direct summand of P . $\square$

The following lemma collects some basic properties of the ideals $\mathfrak{b}^{[1/q]}$.

Lemma 2.4. *Let $\mathfrak{a}$ and $\mathfrak{b}$ be ideals in R and $q = p^e$, $q' = p^{e'}$ where e and e' are positive integers.*

- (1) *If $\mathfrak{a} \subseteq \mathfrak{b}$, then $\mathfrak{a}^{[1/q]} \subseteq \mathfrak{b}^{[1/q]}$.*
- (2) *$(\mathfrak{a} \cap \mathfrak{b})^{[1/q]} \subseteq \mathfrak{a}^{[1/q]} \cap \mathfrak{b}^{[1/q]}$ and $\mathfrak{a}^{[1/q]} + \mathfrak{b}^{[1/q]} = (\mathfrak{a} + \mathfrak{b})^{[1/q]}$.*
- (3) *$(\mathfrak{a} \cdot \mathfrak{b})^{[1/q]} \subseteq \mathfrak{a}^{[1/q]} \cdot \mathfrak{b}^{[1/q]}$.*
- (4) *$(\mathfrak{b}^{[q']})^{[1/q]} = \mathfrak{b}^{[q'/q]} \subseteq (\mathfrak{b}^{[1/q]})^{[q']}$.*
- (5) *$\mathfrak{b}^{[1/qq']} \subseteq (\mathfrak{b}^{[1/q]})^{[1/q']}$.*
- (6) *$\mathfrak{b}^{[1/q]} \subseteq (\mathfrak{b}^{q'})^{[1/qq']}$.*

Proof. Statement (1) is straightforward from definition, and both inclusions " $\subseteq$ " in (2) follow from (1). On the other hand, in order to show that $(\mathfrak{a} + \mathfrak{b})^{[1/q]} \subseteq \mathfrak{a}^{[1/q]} + \mathfrak{b}^{[1/q]}$, it is enough to use the fact that

$$\mathfrak{a} + \mathfrak{b} \subseteq (\mathfrak{a}^{[1/q]})^{[q]} + (\mathfrak{b}^{[1/q]})^{[q]} = (\mathfrak{a}^{[1/q]} + \mathfrak{b}^{[1/q]})^{[q]},$$

and the minimality in the definition of $(\mathfrak{a} + \mathfrak{b})^{[1/q]}$.

For (3), note that since we have $\mathfrak{a} \subseteq (\mathfrak{a}^{[1/q]})^{[q]}$ and $\mathfrak{b} \subseteq (\mathfrak{b}^{[1/q]})^{[q]}$, we deduce

$$\mathfrak{a} \cdot \mathfrak{b} \subseteq (\mathfrak{a}^{[1/q]})^{[q]} \cdot (\mathfrak{b}^{[1/q]})^{[q]} = (\mathfrak{a}^{[1/q]} \cdot \mathfrak{b}^{[1/q]})^{[q]}.$$

The inclusion in (3) then follows from the minimality in the definition of $(\mathfrak{a} \cdot \mathfrak{b})^{[1/q]}$.

Statement (4) is straightforward when $q' \geq q$, so we assume that $q = p^e \geq q' = p^{e'}$. For any ideal J in a regular ring, we have $\mathfrak{b}^{[p^{e'}]} \subseteq J^{[p^e]}$ if and only if $\mathfrak{b} \subseteq J^{[p^{e-e'}]}$. On the other hand, note that since

$$\mathfrak{b} \subseteq (\mathfrak{b}^{[1/p^e]})^{[p^e]} = \left((\mathfrak{b}^{[1/p^e]})^{[p^{e'}]} \right)^{[p^{e-e'}]},$$

we get $\mathfrak{b}^{[1/p^{e-e'}]} \subseteq (\mathfrak{b}^{[1/p^e]})^{[p^{e'}]}$ from the minimality in the definition of $\mathfrak{b}^{[1/p^{e-e'}]}$.

Similarly, for (5) we use the minimality in the definition of $\mathfrak{b}^{[1/qq']}$ and the fact that

$$\mathfrak{b} \subseteq (\mathfrak{b}^{[1/q]})^{[q]} \subseteq \left((\mathfrak{b}^{[1/q]})^{[1/q']} \right)^{[qq']}.$$

Finally, for (6), note that (4) implies $\mathfrak{b}^{[1/q]} = (\mathfrak{b}^{[q']})^{[1/qq']}$, and this is contained in $(\mathfrak{b}^{q'})^{[1/qq']}$ by (1). $\square$

When R is free over R^{p^e} one has the following alternative description of these newly defined ideals. Again, in the case of a principal ideal this was already observed in [AMBL].

Proposition 2.5. *Suppose that R is free over R^q and let $e_1, \dots, e_N$ be a basis of R over R^q . If $h_1, \dots, h_s$ are generators of an ideal $\mathfrak{b}$ of R , and if for every $i = 1, \dots, s$ we write*

$$(2) \quad h_i = \sum_{j=1}^N a_{i,j}^q e_j$$

with $a_{i,j} \in R$, then $\mathfrak{b}^{[1/q]} = (a_{i,j} \mid i \leq s, j \leq N)$.

Note that the proposition implies in particular that the description therein does not depend on the chosen basis of R over R^q or on the generators of $\mathfrak{b}$.

Proof of Proposition 2.5. It follows from (2) that $(h_1, \dots, h_s) \subseteq (a_{i,j}^q \mid i \leq s, j \leq N)$ and therefore $\mathfrak{b} \subseteq (a_{i,j} \mid i, j)^{[q]}$. Hence the inclusion “ $\subseteq$ ” in the proposition follows immediately from the definition of $\mathfrak{b}^{[1/q]}$.

For the reverse inclusion, suppose that $\mathfrak{b} \subseteq J^{[q]}$. If $g_1, \dots, g_m$ generate J , we may write

$$(3) \quad h_i = \sum_{\ell=1}^m b_\ell g_\ell^q$$

for some $b_\ell \in R$. Remembering that R is assumed free over R^q , consider the dual basis $e_1^*, \dots, e_N^*$ for $\text{Hom}_{R^q}(R, R^q)$, so $e_i^*(e_j) = \delta_{i,j}$. It follows from (2) that $e_j^*(h_i) = a_{i,j}^q$. On

the other hand, (3) shows that

$$e_j^*(h_i) = \sum_{\ell} g_{\ell}^q e_j^*(b_{\ell}) \in J^{[q]}.$$

Therefore $a_{i,j} \in J$ for every i and j , which gives $(a_{i,j} \mid i \leq s, j \leq N) \subseteq \mathfrak{b}^{[1/q]}$. $\square$

Remark 2.6. In [AMBL] it is shown that the ideals $\mathfrak{b}^{[1/q]}$ have the following D -module theoretic description. If $D_R^{(e)}$ denotes the subring of the ring of differential operators D_R which are linear over R^q (recall that $q = p^e$), then $(\mathfrak{b}^{[1/q]})^{[q]}$ is equal to the $D_R^{(e)}$ -submodule of R generated by $\mathfrak{b}$. This interesting viewpoint will not be exploited further here. It is however used successfully in our preprint [BMS] to derive rationality and discreteness of F -thresholds of principal ideals in rings of formal power series, for example; thereby generalizing the 2 variable case treated in [HM].

The following lemma shows that the formation of the ideals $\mathfrak{b}^{[1/q]}$ commutes with localization and completion. In particular, in order to compute $\mathfrak{b}^{[1/q]}$ we may always localize so that R is free over R^q and then use Proposition 2.5.

Lemma 2.7. *Let $\mathfrak{b}$ be an ideal in R .*

- (1) *If S is a multiplicative system in R , then $(S^{-1}\mathfrak{b})^{[1/q]} = S^{-1}(\mathfrak{b}^{[1/q]})$.*
- (2) *If R is local and $\widehat{R}$ is the completion of R , then $(\mathfrak{b}\widehat{R})^{[1/q]} = (\mathfrak{b}^{[1/q]})\widehat{R}$.*

Proof. For (1), note that $\mathfrak{b} \subseteq (\mathfrak{b}^{[1/q]})^{[q]}$ implies

$$S^{-1}\mathfrak{b} \subseteq S^{-1}((\mathfrak{b}^{[1/q]})^{[q]}) = (S^{-1}\mathfrak{b}^{[1/q]})^{[q]}.$$

Therefore $(S^{-1}\mathfrak{b})^{[1/q]} \subseteq S^{-1}(\mathfrak{b}^{[1/q]})$.

For the reverse inclusion, write $(S^{-1}\mathfrak{b})^{[1/q]} = S^{-1}J$, for some ideal J such that $(J : s) = J$ for every $s \in S$. Using the flatness of F we see that $(J^{[q]} : s^q) = J^{[q]}$ for every $s \in S$, hence $(J^{[q]} : s) = J^{[q]}$. Since

$$S^{-1}\mathfrak{b} \subseteq (S^{-1}J)^{[q]} = S^{-1}(J^{[q]}),$$

it follows that $\mathfrak{b} \subseteq J^{[q]}$. Therefore $\mathfrak{b}^{[1/q]} \subseteq J$, which gives “ $\supseteq$ ” in (1).

For (2) we will use Proposition 2.5. Since R is local, R is free over R^q . Moreover, one shows as in Example 2.1 that if $e_1, \dots, e_N$ give a basis of R over R^q , then these elements also give a basis of $\widehat{R}$ over $(\widehat{R})^q$, and the assertion in (2) follows from Proposition 2.5. $\square$

2.2. Generalized test ideals. We will now use the ideals $(\mathfrak{a}^r)^{[1/p^e]}$ to define the generalized test ideals of Hara and Yoshida [HY]. In Proposition 2.22 it is shown that our definition coincides with that in [HY].

Lemma 2.8. *Let $\mathfrak{a}$ be an ideal in R . If r, r', e and e' are such that $\frac{r}{p^e} \geq \frac{r'}{p^{e'}}$ and $e' \geq e$, then*

$$(\mathfrak{a}^r)^{[1/p^e]} \subseteq (\mathfrak{a}^{r'})^{[1/p^{e'}]}.$$

Proof. Note that $r' \leq rp^{e'-e}$, hence $\mathfrak{a}^{r'} \supseteq \mathfrak{a}^{rp^{e'-e}}$. It follows from Lemma 2.4 (1) and (6) that

$$(\mathfrak{a}^{r'})^{[1/p^{e'}]} \supseteq (\mathfrak{a}^{rp^{e'-e}})^{[1/p^{e'}]} \supseteq (\mathfrak{a}^r)^{[1/p^e]}.$$

□

Let $\mathfrak{a}$ be an ideal in R and let c be a positive real number. If we denote by $\lceil x \rceil$ the smallest integer greater than or equal to x , then for every e we have $\frac{\lceil cp^e \rceil}{p^e} \geq \frac{\lceil cp^{e+1} \rceil}{p^{e+1}}$. It follows from Lemma 2.8 that

$$(\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]} \subseteq (\mathfrak{a}^{\lceil cp^{e+1} \rceil})^{[1/p^{e+1}]}.$$

Definition 2.9. With notation as above, the generalized test ideal of $\mathfrak{a}$ with exponent c is defined to be

$$\tau(\mathfrak{a}^c) = \bigcup_{e>0} (\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]}.$$

Since R is Noetherian, this union stabilizes after finitely many steps. In particular, the test ideal $\tau(\mathfrak{a}^c)$ is equal to $(\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]}$ for all sufficiently large e .

Remark 2.10. We can write $R = R_1 \times \cdots \times R_m$ where all R_i are (F -finite) regular domains. An ideal $\mathfrak{a}$ in R can be written as $\mathfrak{a} = \mathfrak{a}_1 \times \cdots \times \mathfrak{a}_m$ and it is clear that for every c we have

$$(4) \quad \tau(\mathfrak{a}^c) = \tau(\mathfrak{a}_1^c) \times \cdots \times \tau(\mathfrak{a}_m^c).$$

This allows us to assume that R is a domain whenever this is convenient.

We make the convention that if R is a domain, then $\tau(\mathfrak{a}^0) = R$ unless $\mathfrak{a} = (0)$, in which case $\tau(\mathfrak{a}^0) = (0)$. When R is not necessarily a domain, we define $\tau(\mathfrak{a}^0)$ such that the decomposition (4) holds also when $c = 0$.

Proposition 2.11. *Let $\mathfrak{a}$ and $\mathfrak{b}$ be ideals of R .*

- (1) *If $c_1 < c_2$, then $\tau(\mathfrak{a}^{c_2}) \subseteq \tau(\mathfrak{a}^{c_1})$.*
- (2) *If $\mathfrak{a} \subseteq \mathfrak{b}$, then $\tau(\mathfrak{a}^c) \subseteq \tau(\mathfrak{b}^c)$.*
- (3) *$\tau((\mathfrak{a} \cap \mathfrak{b})^c) \subseteq \tau(\mathfrak{a}^c) \cap \tau(\mathfrak{b}^c)$ and $\tau(\mathfrak{a}^c) + \tau(\mathfrak{b}^c) \subseteq \tau((\mathfrak{a} + \mathfrak{b})^c)$.*
- (4) *$\tau((\mathfrak{a} \cdot \mathfrak{b})^c) \subseteq \tau(\mathfrak{a}^c) \cdot \tau(\mathfrak{b}^c)$.*

Proof. Lemma 2.4(1) implies that $(\mathfrak{a}^{\lceil c_2 p^e \rceil})^{[1/p^e]} \subseteq (\mathfrak{a}^{\lceil c_1 p^e \rceil})^{[1/p^e]}$. By taking $e \gg 0$, we get the assertion in (1). The other assertions follow similarly, by taking the limit in the corresponding assertions from Lemma 2.4. □

Remark 2.12. The inclusion in (4) above is the analog of the Subadditivity Theorem for multiplier ideals in characteristic zero (see [Laz], Theorem 9.5.20). See also Theorem 6.10 in [HY] for a different approach.

A direct application of Lemma 2.7 (for $\mathfrak{b} = \mathfrak{a}^{\lceil cp^e \rceil}$ with $e \gg 0$) shows that the formation of test ideals commutes with localization and completion (see also [HT]).

Proposition 2.13. *Let $\mathfrak{a}$ be an ideal in R and c a non-negative real number.*

- (1) *If S is a multiplicative system in R , then $\tau((S^{-1}\mathfrak{a})^c) = S^{-1}\tau(\mathfrak{a}^c)$.*
- (2) *If R is local and $\widehat{R}$ is the completion of R , then $\tau((\mathfrak{a}\widehat{R})^c) = \tau(\mathfrak{a}^c)\widehat{R}$.*

We now show that the family of test ideals $\tau(\mathfrak{a}^c)$ of a fixed ideal $\mathfrak{a}$ is right continuous in c .

Proposition 2.14. *If $\mathfrak{a}$ is an ideal in R and c is a non-negative real number, then there exists $\varepsilon > 0$ such that $\tau(\mathfrak{a}^c) = (\mathfrak{a}^r)^{\lceil 1/p^e \rceil}$ whenever $c < \frac{r}{p^e} < c + \varepsilon$. That is, $\tau(\mathfrak{a}^c) = \tau(\mathfrak{a}^{c'})$ where c' is a rational number of the form $\frac{r}{p^e}$ which approximates c from above sufficiently well.*

Proof. We show first that there is $\varepsilon > 0$ and an ideal I in R such that $(\mathfrak{a}^r)^{\lceil 1/p^e \rceil} = I$ whenever $c < \frac{r}{p^e} < c + \varepsilon$. Indeed, otherwise we can find r_m and e_m for $m \geq 1$ such that $\frac{r_m}{p^{e_m}}$ form a strictly decreasing sequence converging to c and $(\mathfrak{a}^{r_m})^{\lceil 1/p^{e_m} \rceil} \neq (\mathfrak{a}^{r_{m+1}})^{\lceil 1/p^{e_{m+1}} \rceil}$ for every m . After replacing this sequence by a subsequence we may assume that $e_m \leq e_{m+1}$ for every m . By Lemma 2.8, we have $(\mathfrak{a}^{r_m})^{\lceil 1/p^{e_m} \rceil} \subsetneq (\mathfrak{a}^{r_{m+1}})^{\lceil 1/p^{e_{m+1}} \rceil}$ for every m . Since this sequence of ideals does not stabilize, we contradict the fact that R is Noetherian. Therefore we can find an ideal I as claimed.

We show now that $I = \tau(\mathfrak{a}^c)$. By Remark 2.10 we may assume that R is a domain. If $\mathfrak{a} = (0)$, then $(\mathfrak{a}^r)^{\lceil 1/p^e \rceil} = (0)$ for every r and e , and our assertion is trivial. We assume henceforth that $\mathfrak{a} \neq (0)$. Let e be large enough such that $\tau(\mathfrak{a}^c) = (\mathfrak{a}^{\lceil cp^e \rceil})^{\lceil 1/p^e \rceil}$ and $\frac{\lceil cp^e \rceil}{p^e} < c + \varepsilon$. If cp^e is not an integer, then $\frac{\lceil cp^e \rceil}{p^e} > c$, and we get $I = \tau(\mathfrak{a}^c)$.

Suppose now that cp^e is an integer. After possibly replacing e by a larger value, we may assume that $c + \frac{1}{p^e} < c + \varepsilon$, hence $I = (\mathfrak{a}^{cp^e+1})^{\lceil 1/p^e \rceil} \subseteq (\mathfrak{a}^{cp^e})^{\lceil 1/p^e \rceil} = \tau(\mathfrak{a}^c)$. For the reverse inclusion we need to show that $\mathfrak{a}^{cp^e} \subseteq I^{\lceil p^e \rceil}$. Let $u \in \mathfrak{a}^{cp^e}$. If $e' \geq e$, then $\frac{cp^{e'}+1}{p^{e'}} < c + \varepsilon$, hence $\mathfrak{a}^{cp^{e'}+1} \subseteq I^{\lceil p^{e'} \rceil}$. We deduce that if v is a nonzero element in $\mathfrak{a}$, then for every $e' \gg e$ we have $vu^{p^{e'}-e} \in (I^{\lceil p^{e'} \rceil})^{\lceil p^{e'}-e \rceil}$. This says that u is in the tight closure of the ideal $I^{\lceil p^e \rceil}$, which is equal to $I^{\lceil p^e \rceil}$ since R is a regular ring (see [HH]). Therefore we conclude that $u \in I^{\lceil p^e \rceil}$, and the proof is complete. [To avoid the explicit appearance of tight closure one could first reduce to the local case and use the freeness of R over $R^{p^{e'}-e}$ to find, for $e' \gg e$, a splitting $\varphi : R \rightarrow R$ of the $(e' - e)^{\text{th}}$ iterate of the

Frobenius $F^{e'-e} : R \rightarrow R$ such that $\varphi(v) = 1$. Applying φ to an equation witnessing the membership $vu^{p^{e'-e}} \in (I^{[p^e]})^{[p^{e'-e}]}$, it follows that $u \in I^{[p^e]}$. $\square$

Corollary 2.15. *If m is a positive integer, then for every $c \in \mathbb{R}_{\geq 0}$ we have*

$$\tau((\mathfrak{a}^m)^c) = \tau(\mathfrak{a}^{cm}).$$

Proof. It is clear that $((\mathfrak{a}^m)^r)^{[1/p^e]} = (\mathfrak{a}^{rm})^{[1/p^e]}$ for every positive integer r and e . Let e be large enough such that

$$\tau((\mathfrak{a}^m)^c) = (\mathfrak{a}^{\lceil cp^e \rceil m})^{[1/p^e]}, \text{ and } \tau(\mathfrak{a}^{cm}) = (\mathfrak{a}^{\lceil cmp^e \rceil})^{[1/p^e]}.$$

If, for some e as above, we have $cp^e \in \mathbb{Z}$, then our assertion is clear. If this is not the case, then for $e \gg 0$, we have $\frac{\lceil cp^e \rceil m}{p^e}$ larger than cm , but arbitrarily close to cm as e gets large, and the statement follows from Proposition 2.14. $\square$

Corollary 2.16. *For every ideal $\mathfrak{a}$ in R and every non-negative real number c , there is $\varepsilon > 0$ such that $\tau(\mathfrak{a}^c) = \tau(\mathfrak{a}^{c'})$ for every $c' \in [c, c + \varepsilon)$.*

Proof. It is clear that we may take ε as given by Proposition 2.14. $\square$

Definition 2.17. A positive real number c is an *F -jumping exponent* for $\mathfrak{a}$ if $\tau(\mathfrak{a}^c) \neq \tau(\mathfrak{a}^{c-\varepsilon})$ for every positive ε .

Unless explicitly mentioned otherwise, we make the convention that 0 is also an F -jumping exponent. We will study the basic properties of these numbers in the next section.

Remark 2.18. Suppose that K/k is an extension of perfect fields, and consider the ring extension $R = k[x_1, \dots, x_n] \subseteq S = K[x_1, \dots, x_n]$. If $\mathfrak{a}$ is an ideal in R , then $\tau(\mathfrak{a}^c) \cdot S = \tau((\mathfrak{a} \cdot S)^c)$. Indeed, the monomials of degree at most $p^e - 1$ in each variable give a basis of both R and S over R^{p^e} and S^{p^e} , respectively. It follows from Proposition 2.5 that for every r and e we have $(\mathfrak{a}^r)^{[1/p^e]} \cdot S = ((\mathfrak{a}S)^r)^{[1/p^e]}$. In particular, we see that $\mathfrak{a}$ and $\mathfrak{a} \cdot S$ have the same F -jumping exponents.

2.3. The connection with the Hara-Yoshida definition. We show now that the ideals $\tau(\mathfrak{a}^c)$ we have defined coincide with the $\mathfrak{a}^c$ -test ideals introduced in [HY], temporarily denoted by $\tau'(\mathfrak{a}^c)$. To state Hara and Yoshida's definition, we need to recall their notion of $\mathfrak{a}^c$ -tight closure.

For every $e \geq 1$, let R^e denote the $R - R$ -bimodule on R , with the left R -module structure being the usual one, while the right one being induced by $F^e : R \rightarrow R$. For any inclusion $N \subset M$ of R -modules, there is an induced map of left R -modules

$$R^e \otimes N \rightarrow R^e \otimes M$$

whose image we denote by $N_M^{[p^e]}$. An element $m \in M$ is said to be in the $\mathfrak{a}^c$ -tight closure of N in M if there exists $u \in R$, not in any minimal prime, such that

$$\text{image } (u\mathfrak{a}^{[cp^e]} \otimes m) \in N_M^{[p^e]}$$

for all $e \gg 0$. The set of all such elements forms a submodule $N_M^{*\mathfrak{a}^c}$ of M .

We are mainly interested in the case where R is regular, N is zero, and

$$(5) \quad M = E := \bigoplus_{\mathfrak{m} \in \max \text{spec } R} E_R(R/\mathfrak{m})$$

is the direct sum, over all the maximal ideals of R , of the injective hulls at the corresponding residue fields. With this notation, Hara and Yoshida's definition can be stated as follows:

Definition 2.19. Let R be a regular ring of prime characteristic p , $\mathfrak{a}$ an ideal of R and c any positive real number. The $\mathfrak{a}^c$ -test ideal, denoted $\tau'(\mathfrak{a}^c)$, is the annihilator in R of the R -module $0_E^{*\mathfrak{a}^c}$, where E is as above in (5).

Remark 2.20. More generally, Hara and Yoshida define the ideals $\tau'(\mathfrak{a}^c)$ even when R is not regular as the annihilator of the R -module $\bigcup_M 0_M^{*\mathfrak{a}^c}$, where the union is taken over all finitely generated submodules of E .¹ In the regular case, however, it is easy to check that $\bigcup_M 0_M^{*\mathfrak{a}^c} = 0_E^{*\mathfrak{a}^c}$, since the flatness of Frobenius implies $R^e \otimes M \subseteq R^e \otimes E$ for all submodules M of E .

Remark 2.21. Hara and Yoshida's definition recovers the classical definition of the test ideal, as introduced by Hochster and Huneke in [HH], by taking $\mathfrak{a}$ to be the unit ideal, and c arbitrary; see [HH, Prop 8.23]. We caution the reader, however, that the name “test ideal” is misleading if $\mathfrak{a} \neq (1)$, because the elements of $\tau'(\mathfrak{a}^c)$ are *not the test elements* for $\mathfrak{a}^c$ -tight closure. See [HY], Definition 1.6 and Remark 1.10.

The next proposition says that these two notions of test ideals agree.

Proposition 2.22. *If $\mathfrak{a}$ is an ideal in a regular F -finite ring R , then for every non-negative real number c we have $\tau(\mathfrak{a}^c) = \tau'(\mathfrak{a}^c)$.*

Before proving this, we recall a few well-known facts. First, if $(R, \mathfrak{m})$ is a regular local ring of dimension n , then the injective hull of the residue field $E = E_R(R/\mathfrak{m})$ can be identified with the local cohomology module $H_{\mathfrak{m}}^n(R)$. The Frobenius morphism on R induces a Frobenius action on $H_{\mathfrak{m}}^n(R)$ and hence also on E . This action, also denoted by F , can be described as follows. If $x_1, \dots, x_n$ form a minimal system of generators of $\mathfrak{m}$ then

$$E \simeq R_{x_1 \dots x_n} / \sum_{i=1}^n R_{x_1 \dots \widehat{x_i} \dots x_n}.$$

¹In the parlance of [HH], this union could be called the *finitistic $\mathfrak{a}^c$ -tight closure* of zero in E .

Under this isomorphism, the Frobenius morphism on E is induced by the Frobenius morphism on $R_{x_1 \cdots x_n}$. In particular, an element $\eta \in E$ represented by the fraction $\frac{w}{(x_1 \cdots x_n)^N}$ will be sent under Frobenius to the element $\eta^{p^e} \in E$ represented by the fraction

$$\frac{w^{p^e}}{(x_1 \cdots x_n)^{Np^e}}.$$

Also using this representation of $\eta \in E$, we have $\text{Ann}_R \eta = (x_1^N, \dots, x_n^N) :_R w$. In particular, for all $\eta \in E$ we have

$$(6) \quad \text{Ann}_R(\eta^{p^e}) = (\text{Ann}_R \eta)^{[p^e]},$$

for all $e \geq 1$. Indeed, $\text{Ann}_R \eta^{p^e} = (x_1^{p^e N}, \dots, x_n^{p^e N}) :_R w^{p^e} = ((x_1^N, \dots, x_n^N) :_R w)^{[p^e]}$, with the last equality following from the flatness of Frobenius.

The Frobenius action on E induces an R -linear map $\nu^e : R^e \otimes_R E \rightarrow E$ given by $\nu^e(a \otimes \eta) = a\eta^{p^e}$. In the case that R is regular, this map is an isomorphism for all $e \geq 1$. (See, for example, [Lyu], Example 1.2 or [Bli], p.20.) Thus we can conveniently compute the $\mathfrak{a}^c$ -tight closure of zero in E as the set of all $\eta \in E$ for which there exists $u \in R$ not in any minimal prime such that

$$(7) \quad u\mathfrak{a}^{[cp^e]} \eta^{p^e} = 0, \quad \text{for all } e \gg 0.$$

We now prove two lemmas, both well-known to practitioners of tight closure, whose proofs we include to keep the paper self-contained. In the language of tight closure, the first says that, for a regular local ring, zero is tightly closure in the injective hull of the residue field.

Lemma 2.23. *With the above notation, let u be a non-zero element in a regular local ring $(R, \mathfrak{m})$. If $\eta \in E = E_R(R/\mathfrak{m})$ satisfies $u\eta^{p^e} = 0$ for every $e \gg 0$, then $\eta = 0$.*

Proof. The hypothesis implies that $u \in \text{Ann}_R(\eta^{p^e})$ for all $e \gg 0$. But then by (6), we see that $u \in (\text{Ann}_R(\eta))^{[p^e]}$ for every $e \gg 0$. By Nakayama's Lemma, since u is nonzero, it follows that $\text{Ann}_R(\eta) = R$. $\square$

The second lemma is essentially a special case of the statement that 1 is a $\mathfrak{a}^c$ -test element in a regular local ring (see [HY, Theorem 1.7]).

Lemma 2.24. *If R is a regular local ring, then*

$$0_E^{*\mathfrak{a}^c} = \{\eta \in E \mid \mathfrak{a}^{[cp^e]} \eta^{p^e} = 0 \text{ for all } e \geq 1\}.$$

Proof. Clearly $\supseteq$ holds, so consider any $\eta \in 0_E^{*\mathfrak{a}^c}$. We know that there is a non-zero element $u \in R$ such that $u\mathfrak{a}^{[cp^e]} \eta^{p^e} = 0$.

Now, fix any $e' \geq 1$, and any $h \in \mathfrak{a}^{[cp^{e'}]}$. For all $e \geq 1$, we have

$$h^{p^e} \in (\mathfrak{a}^{[cp^{e'}]})^{p^e} \subseteq \mathfrak{a}^{[cp^{e'}+e]}.$$

Thus

$$u(h\eta^{p^{e'}})^{p^e} = uh^{p^e}\eta^{p^{e'+e}} \in u\mathfrak{a}^{\lceil cp^{e'+e} \rceil} \eta^{p^{e'+e}},$$

which is zero for $e \gg 0$ by hypothesis. By Lemma 2.23, it follows that $h\eta^{p^{e'}} = 0$ for all $e' \geq 1$. Since h was an arbitrary element of $\mathfrak{a}^{\lceil cp^{e'} \rceil}$, we conclude that $\mathfrak{a}^{\lceil cp^{e'} \rceil} \eta^{p^{e'}} = 0$ for all $e' \geq 1$. The proof is complete. $\square$

Proof of Proposition 2.22. The proof reduces immediately to the case R is local. Indeed, since $E_R(R/\mathfrak{m})$ is already a $R_{\mathfrak{m}}$ -module, the $\mathfrak{a}^c$ -tight closure of zero in $E_R(R/\mathfrak{m})$ is the same as the $(\mathfrak{a}R_{\mathfrak{m}})^c$ -tight closure of zero in $E_{R_{\mathfrak{m}}}(R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}})$, whence

$$(8) \quad \tau'(\mathfrak{a}^c) = \cap_{\mathfrak{m}} \{f \in R \mid f/1 \in \tau((\mathfrak{a}R_{\mathfrak{m}})^c)\}.$$

This implies that $\tau'(\mathfrak{a}^c)$ commutes with localization at maximal ideals, whereas Lemma 2.7 implies that $\tau(\mathfrak{a}^c)$ does.

Now, we assume $(R, \mathfrak{m})$ is a regular local ring. To show that the ideals $\tau'(\mathfrak{a}^c)$ and $\tau(\mathfrak{a}^c)$ coincide, it is enough, by Matlis duality, to show that their annihilators in E coincide. By definition (and Matlis duality), the annihilator of $\tau'(\mathfrak{a}^c)$ in E is precisely $0_E^{*\mathfrak{a}^c}$. It remains to understand the annihilator in E of $\tau(\mathfrak{a}^c)$.

By definition, $\tau(\mathfrak{a}^c) = (\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]}$ for all $e \gg 0$. So an element η represented by a fraction $\frac{w}{(x_1 \dots x_n)^N}$ is annihilated by $\tau(\mathfrak{a}^c)$ if and only if

$$(\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]} \subseteq (x_1^N, \dots, x_n^N) : w.$$

By definition of $\mathfrak{b}^{[1/q]}$, this holds if and only if

$$\mathfrak{a}^{\lceil cp^e \rceil} \subseteq ((x_1^N, \dots, x_n^N) : w)^{[p^e]} = (x_1^{Np^e}, \dots, x_n^{Np^e}) : w^{p^e},$$

which in turn exactly says that $\mathfrak{a}^{\lceil cp^e \rceil} \eta^{p^e} = 0$ for all $e \gg 0$. We conclude using Lemma 2.24 that the annihilator of $\tau(\mathfrak{a}^c)$ in E is precisely $0_E^{*\mathfrak{a}^c}$. This completes the proof. $\square$

2.4. Skoda's Theorem. For future reference, we include the following characteristic p version of Skoda's Theorem, due to Hara and Takagi [HT]. Since the result in *loc. cit.* is not stated in the generality we will need, we include a proof for the benefit of the reader.

Proposition 2.25. *If $\mathfrak{a}$ is an ideal generated by m elements, then for every $c \geq m$ we have*

$$\tau(\mathfrak{a}^c) = \mathfrak{a} \cdot \tau(\mathfrak{a}^{c-1}).$$

Proof. If e is large enough, then $\tau(\mathfrak{a}^c) = (\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]}$ and $\tau(\mathfrak{a}^{c-1}) = (\mathfrak{a}^{\lceil cp^e \rceil - p^e})^{[1/p^e]}$. Therefore it is enough to show that for every $r \geq mp^e$ we have

$$(\mathfrak{a}^r)^{[1/p^e]} = \mathfrak{a} \cdot (\mathfrak{a}^{r-p^e})^{[1/p^e]}.$$

The inclusion $\mathfrak{a} \cdot (\mathfrak{a}^{r-p^e})^{[1/p^e]} \subseteq (\mathfrak{a}^r)^{[1/p^e]}$ holds in fact for every $r \geq p^e$. Indeed, this says that $(\mathfrak{a}^{r-p^e})^{[1/p^e]} \subseteq ((\mathfrak{a}^r)^{[1/p^e]} : \mathfrak{a})$, which is equivalent with

$$\mathfrak{a}^{r-p^e} \subseteq ((\mathfrak{a}^r)^{[1/p^e]} : \mathfrak{a})^{[p^e]} = \left(((\mathfrak{a}^r)^{[1/p^e]})^{[p^e]} : \mathfrak{a}^{[p^e]} \right).$$

This holds since $\mathfrak{a}^{r-p^e} \cdot \mathfrak{a}^{[p^e]} \subseteq \mathfrak{a}^r \subseteq ((\mathfrak{a}^r)^{[1/p^e]})^{[p^e]}$.

Suppose now that $\mathfrak{a} = (h_1, \dots, h_m)$. In order to prove the reverse inclusion, note that if $r \geq m(p^e - 1) + 1$, then in the product of r of the h_i , at least one of these appears p^e times. Therefore $\mathfrak{a}^r = \mathfrak{a}^{[p^e]} \cdot \mathfrak{a}^{r-p^e}$. We deduce that

$$\mathfrak{a}^r \subseteq \mathfrak{a}^{[p^e]} \cdot \mathfrak{a}^{r-p^e} \subseteq \mathfrak{a}^{[p^e]} \cdot \left((\mathfrak{a}^{r-p^e})^{[1/p^e]} \right)^{[p^e]} = (\mathfrak{a} \cdot (\mathfrak{a}^{r-p^e})^{[1/p^e]})^{[p^e]},$$

which implies $(\mathfrak{a}^r)^{[1/p^e]} \subseteq \mathfrak{a} \cdot (\mathfrak{a}^{r-p^e})^{[1/p^e]}$. $\square$

Remark 2.26. In Proposition 2.25, rather than taking m to be the minimal number of generators for $\mathfrak{a}$, we can take m to be the *analytic spread* of $\mathfrak{a}$, that is, the minimal number of generators for any subideal with the same integral closure as $\mathfrak{a}$. Indeed, this is immediate from the following lemma, showing that the test ideals of $\mathfrak{a}$ depend only on its integral closure (see also [HY] and [HT]).

Lemma 2.27. *If $\bar{\mathfrak{a}}$ denotes the integral closure of $\mathfrak{a}$, then $\tau(\mathfrak{a}^c) = \tau(\bar{\mathfrak{a}}^c)$ for every c .*

Proof. The inclusion $\tau(\mathfrak{a}^c) \subseteq \tau(\bar{\mathfrak{a}}^c)$ is immediate from Proposition 2.11 (2). For the reverse inclusion, note that by usual properties of integral closure, there exists m such that $\bar{\mathfrak{a}}^{m+\ell} \subseteq \mathfrak{a}^\ell$ for every ℓ . Corollary 2.16 gives $c' > c$ such that $\tau(\mathfrak{a}^c) = \tau(\mathfrak{a}^{c'})$ and $\tau(\bar{\mathfrak{a}}^c) = \tau(\bar{\mathfrak{a}}^{c'})$.

Using Corollary 2.15, we see that

$$\tau(\bar{\mathfrak{a}}^{c'}) = \tau((\bar{\mathfrak{a}}^{m+\ell})^{\frac{c'}{m+\ell}}) \subseteq \tau((\mathfrak{a}^\ell)^{\frac{c'}{m+\ell}}) = \tau(\mathfrak{a}^{c' - \frac{c'm}{m+\ell}}).$$

If $\ell \gg 0$, then $c < c' - \frac{c'm}{m+\ell} < c'$, hence by our choice of c' we get $\tau(\bar{\mathfrak{a}}^c) \subseteq \tau(\mathfrak{a}^c)$. $\square$

Remark 2.28. If $\varphi: R \rightarrow S$ is a morphism of regular, F -finite rings of positive characteristic, and if $\mathfrak{b}$ is an ideal in R , then $(\mathfrak{b} \cdot S)^{[1/p^e]} \subseteq \mathfrak{b}^{[1/p^e]} \cdot S$ for every e . Indeed, since $\mathfrak{b} \subseteq (\mathfrak{b}^{[1/p^e]})^{[p^e]}$ we have $\mathfrak{b} \cdot S \subseteq (\mathfrak{b}^{[1/p^e]} \cdot S)^{[p^e]}$.

We deduce that for every non-negative c we have $\tau((\mathfrak{a} \cdot S)^c) \subseteq \tau(\mathfrak{a}^c) \cdot S$. This is an analogue of the Restriction Theorem for multiplier ideals in characteristic zero (see [Laz], Examples 9.5.4 and 9.5.8). For a different argument in a more general (characteristic p) framework, see [HY], Theorems 4.1 and 6.10.

2.5. F -jumping exponents and F -thresholds. In [MTW] F -jumping exponents of an ideal $\mathfrak{a}$ were described as F -thresholds. Since the statements and the proofs in *loc. cit.* were given in the local case, we review them here for the reader's convenience.

Let $\mathfrak{a}$ be an ideal in R . For a fixed ideal J in R such that $\mathfrak{a} \subseteq \text{rad}(J)$ and for an integer $e > 0$ we define $\nu_{\mathfrak{a}}^J(p^e)$ to be the largest non-negative integer r such that $\mathfrak{a}^r \not\subseteq J^{[p^e]}$ (if there is no such r , then we put $\nu_{\mathfrak{a}}^J(p^e) = 0$). If $\mathfrak{a}^r \not\subseteq J^{[p^e]}$, then $\mathfrak{a}^{pr} \not\subseteq J^{[p^{e+1}]}$. Indeed, otherwise we get $(\mathfrak{a}^r)^{[p]} \subseteq J^{[p^{e+1}]}$, hence $\mathfrak{a}^r \subseteq J^{[p^e]}$, a contradiction. Therefore

$$\frac{\nu_{\mathfrak{a}}^J(p^e)}{p^e} \leq \frac{\nu_{\mathfrak{a}}^J(p^{e+1})}{p^{e+1}},$$

hence we may define the F -threshold of $\mathfrak{a}$ with respect to J as

$$c^J(\mathfrak{a}) := \lim_{e \rightarrow \infty} \frac{\nu_{\mathfrak{a}}^J(p^e)}{p^e} = \sup_{e \geq 1} \frac{\nu_{\mathfrak{a}}^J(p^e)}{p^e}.$$

Note that if $\mathfrak{a}$ is generated by s elements, then $\mathfrak{a}^{s(p^e-1)+1} \subseteq \mathfrak{a}^{[p^e]}$. If $\mathfrak{a}^\ell \subseteq J$, then $\nu_{\mathfrak{a}}^J(p^e) \leq \ell(s(p^e - 1) + 1) - 1$ for every e . Therefore $c^J(\mathfrak{a}) \leq s\ell$, in particular $c^J(\mathfrak{a})$ is finite.

The following proposition relates the F -thresholds with the generalized test ideals of $\mathfrak{a}$.

Proposition 2.29. *Let $\mathfrak{a}$ be an ideal in R .*

(1) *If J is an ideal in R such that $\mathfrak{a} \subseteq \text{rad}(J)$, then*

$$\tau(\mathfrak{a}^{c^J(\mathfrak{a})}) \subseteq J.$$

(2) *If c is a non-negative real number, then $\mathfrak{a} \subseteq \text{rad}(\tau(\mathfrak{a}^c))$ and*

$$c^{\tau(\mathfrak{a}^c)}(\mathfrak{a}) \leq c.$$

Proof. For (1), note that by Corollary 2.16 there is $c' > c^J(\mathfrak{a})$ such that $I := \tau(\mathfrak{a}^{c^J(\mathfrak{a})}) = \tau(\mathfrak{a}^{c'})$. Suppose now that $e \gg 0$, so $\tau(\mathfrak{a}^{c'}) = (\mathfrak{a}^{\lceil c'p^e \rceil})^{[1/p^e]}$.

Since $c' > c^J(\mathfrak{a})$ and e is large enough, we have $\lceil c'p^e \rceil \geq \nu_{\mathfrak{a}}^J(p^e) + 1$, hence

$$\mathfrak{a}^{\lceil c'p^e \rceil} \subseteq J^{[p^e]}.$$

This implies that $I \subseteq J$, as required.

For (2), let e be large enough, so $\tau(\mathfrak{a}^c) = (\mathfrak{a}^{\lceil cp^e \rceil})^{[1/p^e]}$. By definition, we have $\mathfrak{a}^{\lceil cp^e \rceil} \subseteq \tau(\mathfrak{a}^c)^{[p^e]}$, which implies

$$\nu_{\mathfrak{a}}^{\tau(\mathfrak{a}^c)}(p^e) \leq \lceil cp^e \rceil - 1.$$

Dividing by p^e and letting e go to infinity, we get the required inequality. $\square$

Corollary 2.30. *For every ideal $\mathfrak{a}$ in R , the set of F -jumping exponents for $\mathfrak{a}$ is equal to the set of F -thresholds of $\mathfrak{a}$ (as we range over all possible ideals J).*

Proof. We show first that if α is an F -jumping exponent for $\mathfrak{a}$, then $\alpha = c^J(\mathfrak{a})$ for $J = \tau(\mathfrak{a}^\alpha)$. Indeed, Proposition 2.29(2) gives $c^J(\mathfrak{a}) \leq \alpha$. Therefore, by Proposition 2.11(1), $J = \tau(\mathfrak{a}^\alpha) \subseteq \tau(\mathfrak{a}^{c^J(\mathfrak{a})})$. We also have the reverse inclusion by Proposition 2.29(1). Since α is an F -jumping exponent, we must have $\alpha = c^J(\mathfrak{a})$.

Suppose now that $\alpha = c^J(\mathfrak{a})$ for some J whose radical contains $\mathfrak{a}$. We need to show that α is a jumping exponent. If this is not the case, then there is $\alpha' < \alpha$ such that $\tau(\mathfrak{a}^\alpha) = \tau(\mathfrak{a}^{\alpha'})$. Using Proposition 2.29(1) we get $\tau(\mathfrak{a}^{\alpha'}) \subseteq J$. If e is large enough, then $\tau(\mathfrak{a}^{\alpha'}) = (\mathfrak{a}^{\lceil \alpha' p^e \rceil})^{[1/p^e]}$. Therefore $\mathfrak{a}^{\lceil \alpha' p^e \rceil} \subseteq J^{[p^e]}$, hence $\nu^J(\mathfrak{a}) \leq \lceil \alpha' p^e \rceil - 1$. Dividing by p^e and letting e go to infinity, we get $c^J(\mathfrak{a}) \leq \alpha'$, a contradiction. This completes the proof of the corollary. $\square$

2.6. Mixed generalized test ideals. Using our approach it is easy to define also "mixed" test ideals, as in [HY] and [HT]. For example, suppose that $\mathfrak{a}_1, \dots, \mathfrak{a}_r$ are ideals in the regular F -finite ring R , and let $c_1, \dots, c_r \in \mathbb{R}_+$. For every $e \geq 1$, consider

$$I_e := \left(\mathfrak{a}_1^{\lceil c_1 p^e \rceil} \cdot \dots \cdot \mathfrak{a}_r^{\lceil c_r p^e \rceil} \right)^{[1/p^e]}.$$

As before, one can check easily that $I_e \subseteq I_{e+1}$ for every e , and since R is Noetherian, this sequence of ideals stabilizes. Its limit is the test ideal $\tau(\mathfrak{a}_1^{c_1} \cdot \dots \cdot \mathfrak{a}_r^{c_r})$. It is straightforward to check that the basic properties we have discussed so far generalize to this setting. For example, the Subadditivity Formula from Proposition 2.11 can be generalized with the same proof to the following statement: if $\mathfrak{a}_1, \dots, \mathfrak{a}_r$ are ideals in R , and if $\lambda_1, \dots, \lambda_r \in \mathbb{R}_+$, then

$$\tau(\mathfrak{a}_1^{\lambda_1} \cdot \dots \cdot \mathfrak{a}_r^{\lambda_r}) \subseteq \tau(\mathfrak{a}_1^{\lambda_1}) \cdot \dots \cdot \tau(\mathfrak{a}_r^{\lambda_r})$$

(see also [HY], Theorem 6.10). Similarly, the Summation Formula due to Takagi (see [Tak1], Theorem 3.1) can be easily proved using our definition.

3. DISCRETENESS AND RATIONALITY

In this section we prove our main result.

Theorem 3.1. *Let k be a field of characteristic $p > 0$ and let R be a regular F -finite ring, essentially of finite type over k . Suppose that $\mathfrak{a}$ is an ideal in R .*

- (1) *The set of F -jumping exponents of $\mathfrak{a}$ is discrete (in every finite interval there are only finitely many such numbers).*
- (2) *Every F -jumping exponent of $\mathfrak{a}$ is a rational number.*

We will reduce the proof of the theorem to the case $R = k[x_1, \dots, x_n]$. We start with some preliminary results. The first proposition, of independent interest, gives an effective bound for the degrees of the generators of the ideals $\tau(\mathfrak{a}^c)$ in terms of the degrees of the generators of $\mathfrak{a}$. It is our main ingredient for the proof of the theorem in

the polynomial ring case. For a real number t we will denote by $\lfloor t \rfloor$ the largest integer less than or equal to t , and by $\{t\}$ the fractional part $t - \lfloor t \rfloor$.

Proposition 3.2. *Let $\mathfrak{a}$ be an ideal in the polynomial ring $k[x_1, \dots, x_n]$, where k is a field of characteristic p such that $[k : k^p] < \infty$. If $\mathfrak{a}$ can be generated by polynomials of degree at most d , then for every non-negative real number c , the ideal $\tau(\mathfrak{a}^c)$ can be generated by polynomials of degree at most $\lfloor cd \rfloor$.*

Proof. Fix first r and e . The ideal $\mathfrak{a}^r$ is generated by polynomials of degree at most rd . Choose such generators $h_1, \dots, h_s$ for $\mathfrak{a}^r$.

Let $b_1, \dots, b_m$ be a basis of k over k^{p^e} , and consider the basis of $R = k[x_1, \dots, x_n]$ over R^{p^e} given by

$$\{b_i x^u \mid i \leq m, u \in \mathbb{N}^n, 0 \leq u_j \leq p^e - 1 \text{ for all } j\}$$

(if $u = (u_1, \dots, u_n)$, then we put $x^u = x_1^{u_1} \cdots x_n^{u_n}$). If we write

$$h_j = \sum_{i,u} a_{i,u}^{p^e} b_i x^u,$$

with $a_{i,u} \in R$, then for every i and u we have $\deg(a_{i,u}^{p^e}) \leq \deg(h_j) \leq rd$. It follows from Proposition 2.5 that $(\mathfrak{a}^r)^{[1/p^e]}$ can be generated by polynomials of degree at most $\frac{rd}{p^e}$. But by Proposition 2.14, $\tau(\mathfrak{a}^c) = (\mathfrak{a}^r)^{[1/p^e]}$ for some $\frac{r}{p^e} \geq c$ closely approximating c . Thus for sufficiently close approximations $\frac{r}{p^e}$, $\tau(\mathfrak{a}^c)$ is generated by polynomials of degree at most $\lfloor \frac{rd}{p^e} \rfloor = \lfloor cd \rfloor$. The proof is complete. $\square$

Remark 3.3. As Rob Lazarsfeld points out, the analog of Proposition 3.2 for multiplier ideals can be deduced from a vanishing statement. Indeed, suppose that $\mathfrak{a} \subseteq \mathbb{C}[x_1, \dots, x_n]$ is an ideal generated in degree at most d , and denote by $\mathcal{J}(\mathfrak{a}^c)$ its multiplier ideal. After homogenizing $\mathfrak{a}$ and taking the associated sheaf on $\mathbf{P}^n$, we get a sheaf of ideals $\tilde{\mathfrak{a}}$ whose restriction to $\mathbb{A}^n = (x_0 \neq 0) \subseteq \mathbf{P}^n$ is $\mathfrak{a}$, and such that $\tilde{\mathfrak{a}} \otimes \mathcal{O}_{\mathbf{P}}(d)$ is globally generated. If H denotes the hyperplane class on $\mathbf{P}^n$, then $\lfloor cd \rfloor H \sim K_{\mathbf{P}^n} + (n + 1 + \lfloor cd \rfloor)H$. For $1 \leq i \leq n$ we have $(n + 1 + \lfloor cd \rfloor - i - cd)H$ ample, and therefore

$$H^i(\mathbf{P}^n, \mathcal{J}(\tilde{\mathfrak{a}}^c) \otimes \mathcal{O}_{\mathbf{P}}(\lfloor cd \rfloor - i)) = 0$$

by the Nadel Vanishing Theorem (see [Laz], Corollary 9.4.15). Therefore $\mathcal{J}(\tilde{\mathfrak{a}}^c) \otimes \mathcal{O}_{\mathbf{P}}(\lfloor cd \rfloor)$ is 0-regular in the sense of Castelnuovo-Mumford regularity, hence it is generated by global sections. This implies that $\mathcal{J}(\mathfrak{a}^c)$ is generated in degree at most $\lfloor cd \rfloor$.

Proposition 3.4. *Let $\mathfrak{a}$ be an ideal in a regular, F -finite ring R .*

- (1) *If α is an F -jumping exponent for $\mathfrak{a}$, then $p\alpha$ is an F -jumping exponent, too.*
- (2) *If $\mathfrak{a}$ can be generated by m elements, and if $\alpha > m$ is an F -jumping exponent for $\mathfrak{a}$, then $\alpha - 1$ is an F -jumping exponent, too.*

Remark 3.5. For a stronger statement in (2), we can take m to be any integer at least as large as the analytic spread of $\mathfrak{a}$. (This follows from Remark 2.26.) For example, if R is local with an infinite residue field, we can take $m = \dim R$.

Proof. For (1), note that by Corollary 2.30 there is an ideal J containing $\mathfrak{a}$ in its radical such that $\alpha = c^J(\mathfrak{a})$. It is clear that $\nu_{\mathfrak{a}}^{J[p]}(p^e) = \nu_{\mathfrak{a}}^J(p^{e+1})$, so $c^{J[p]}(\mathfrak{a}) = p \cdot c^J(\mathfrak{a})$. Hence $p\alpha$ is an F -jumping exponent of $\mathfrak{a}$ by Corollary 2.30.

For (2), suppose that $\alpha - 1$ is not an F -jumping exponent. Let $\varepsilon > 0$ be such that $\tau(\mathfrak{a}^{\alpha-1}) = \tau(\mathfrak{a}^{\alpha-1-\varepsilon})$ and $\alpha - \varepsilon > m$. It follows from Proposition 2.25 that $\tau(\mathfrak{a}^\alpha) = \tau(\mathfrak{a}^{\alpha-\varepsilon})$, hence α is not an F -jumping exponent, a contradiction. $\square$

The following proposition relates the generalized test ideals of two different ideals defining the same scheme. For the characteristic zero analogue in the context of multiplier ideals, see Proposition 2.3 in [Mu].

Proposition 3.6. *Let R be a regular, F -finite ring of positive characteristic, and I an ideal in R of pure codimension r , such that $S = R/I$ is regular. If $\mathfrak{a}$ is an ideal in R containing I , then for every non-negative real number c we have*

$$\tau((\mathfrak{a}/I)^c) = \tau(\mathfrak{a}^{c+r}) \cdot S.$$

Proof. By Proposition 2.13, forming generalized test ideals commutes with localization and completion. Therefore it is enough to prove the case where R is local and complete. Since I is generated by part of a regular system of parameters for R , by doing induction on r we see that it is enough to prove the case $r = 1$. Therefore we may assume that $R = k[[x_1, \dots, x_n]]$ and $I = (x_n)$. Note that $[k : k^p] < \infty$, since R is F -finite.

We claim that

$$(9) \quad (\mathfrak{a}^{r+p^e})^{[1/p^e]} \cdot S = ((\mathfrak{a}/(x_n))^{r+1})^{[1/p^e]}$$

for any integer r and all $e \gg 0$. This implies the desired statement. Indeed, if e is large enough, then we have

$$\tau(\mathfrak{a}^{c+1}) \cdot S = (\mathfrak{a}^{[cp^e]+p^e})^{[1/p^e]} \cdot S = ((\mathfrak{a}/(x_n))^{[cp^e]+1})^{[1/p^e]} = \tau((\mathfrak{a}/(x_n))^c).$$

The last equality follows from Proposition 2.14, since $\frac{[cp^e]+1}{p^e}$ is larger than c , but arbitrarily close to c as e gets very large.

We prove now the claim (9), using the description in Proposition 2.5. Let $a_1, \dots, a_m$ be a basis of k over k^{p^e} , and consider the basis of R over R^{p^e} given by

$$\{a_i x^u \mid i \leq m, u = (u_j) \in \mathbb{N}^n, 0 \leq u_j \leq p^e - 1 \text{ for all } j\}.$$

Write $\mathfrak{a} = (x_n) + \mathfrak{b}$, where $\mathfrak{b}$ is generated by power series in $k[[x_1, \dots, x_{n-1}]]$. We have

$$\mathfrak{a}^{r+p^e} = \sum_{i=0}^{r+p^e} x_n^i \mathfrak{b}^{r+p^e-i}.$$

The generators of $(\mathfrak{a}^{r+p^e})^{[1/p^e]}$ that come from writing in the above basis the generators of $x_n^i \mathfrak{b}^{r+p^e-i}$ are divisible by x_n if $i \geq p^e$, hence map to zero in S . On the other hand, the generators coming from $x_n^i \mathfrak{b}^{r+p^e-i}$ for $i \leq p^e - 1$ are the same as the ones obtained from writing the generators of $\mathfrak{b}^{r+p^e-i}$ in the corresponding basis of $k[[x_1, \dots, x_{n-1}]]$ over $k^{p^e}[[x_1^{p^e}, \dots, x_{n-1}^{p^e}]]$. Moreover, it is clear that it is enough to consider only the largest such ideal, namely $\mathfrak{b}^{r+1}$. This shows that $(\mathfrak{a}^{r+p^e})^{[1/p^e]} \cdot S = ((\mathfrak{a}/(x_n))^{r+1})^{[1/p^e]}$, as claimed. $\square$

Corollary 3.7. *If R , S and $\mathfrak{a}$ are as in Proposition 3.6, and if $c > 0$ is a jumping exponent for $\mathfrak{a} \cdot S$, then $c + r$ is a jumping exponent for $\mathfrak{a}$.*

We can give now the proof of our main result.

Proof of Theorem 3.1. For (1), suppose that we have a sequence of F -jumping exponents $\{\alpha_m\}_m$ for $\mathfrak{a}$ having a finite accumulation point α . By Corollary 2.16, we have $\alpha_m < \alpha$ for $m \gg 0$. After replacing this sequence by a subsequence, we may assume that $\alpha_m < \alpha_{m+1}$ for every m .

Let us write $R = R_1 \times \dots \times R_s$, where all R_i are domains. We have $\mathfrak{a} = \mathfrak{a}_1 \times \dots \times \mathfrak{a}_s$, and for every m there is j such that α_m is a jumping exponent for $\mathfrak{a}_j$. After replacing our sequence by a subsequence, we may replace R by some R_j and therefore assume that R is a domain.

By hypothesis, we can write $R \simeq S^{-1}(k[x_1, \dots, x_n]/I)$ for some ideal I and some multiplicative system S . Note that $[k : k^p] < \infty$ (see Example 2.1). Let us write $\mathfrak{a} = S^{-1}(\mathfrak{b}/I)$, for some ideal $\mathfrak{b} \subset k[x_1, \dots, x_n]$ containing I . Note that $S^{-1}I$ is a prime ideal, hence it has pure codimension, say r , in $S^{-1}k[x_1, \dots, x_n]$. It follows from Corollary 3.7 that $r + \alpha$ is an accumulation point for the jumping exponents of $S^{-1}\mathfrak{b}$. Moreover, Proposition 2.13(1) implies that $r + \alpha$ is an accumulation point for the jumping exponents of $\mathfrak{b}$. Therefore, in order to achieve a contradiction we may assume that $R = k[x_1, \dots, x_n]$.

Suppose now that $\mathfrak{a}$ is generated by polynomials of degree at most d . In this case, Proposition 3.2 implies that every $\tau(\mathfrak{a}^{\alpha_m})$ is generated by polynomials of degree at most $\lfloor \alpha d \rfloor$. Since the ideals $\tau(\mathfrak{a}^{\alpha_m})$ form a strictly decreasing sequence of ideals, their k -subvector spaces of polynomials of degree at most $\lfloor \alpha d \rfloor$ form a strictly decreasing sequence of subspaces of the finite dimensional vector space $k[x_1, \dots, x_n]_{\leq \lfloor \alpha d \rfloor}$. This contradiction completes the proof of (1).

Suppose now that $\alpha > 0$ is an F -jumping exponent for $\mathfrak{a}$. Proposition 3.4(1) implies that all $p^e\alpha$ are F -jumping exponents for $\mathfrak{a}$. We may assume that no $p^e\alpha$ is an integer, as otherwise α is clearly rational. Suppose that $\mathfrak{a}$ is generated by m elements and let e_0 be such that $p^{e_0}\alpha > m$. We deduce from Proposition 3.4(2) that $\{p^e\alpha\} + m - 1$ is a jumping exponent for every $e \geq e_0$. Since all these numbers lie in $[m - 1, m)$, it follows from (1) that there are only finitely many such numbers. Therefore we can find $e_1 \neq e_2$ such that $p^{e_1}\alpha - p^{e_2}\alpha$ is an integer. Hence α is a rational number. $\square$

The ideas in the above proof can be used to explicitly estimate the F -jumping exponents of an ideal in a polynomial ring in terms of the degrees of its generators. Since we know that all F -jumping exponents are rational, it is enough to bound their denominators.

Proposition 3.8. *Let $\mathfrak{a} \subseteq k[x_1, \dots, x_n]$ be an ideal generated by m polynomials of degree at most d . If e_0 is such that $p^{e_0} > md$ and $N = \binom{md+n}{n}$, then for every F -jumping exponent α of $\mathfrak{a}$ we have $p^a(p^b - 1)\alpha \in \mathbb{N}$ for some $a \leq e_0 + N$ and $b \leq N$.*

Proof. By Proposition 3.4(2) it is enough to consider the case when $\alpha \leq m$. Since $\tau(\mathfrak{a}^c)$ is generated by polynomials of degree at most md for every $c \leq m$, it follows that we have at most $N = \dim_k k[x_1, \dots, x_n]_{\leq md}$ jumping exponents of $\mathfrak{a}$ in $(0, m]$.

We may assume that for every $e \leq e_0 + N$ we have $p^e\alpha \notin \mathbb{N}$, since otherwise our assertion is clear. In particular, $\alpha > 0$ and therefore $\alpha \geq 1/d$ (note that $\tau(\mathfrak{a}^c) = R$ for $c < 1/d$ by Proposition 3.2). Consider now e , with $e_0 \leq e \leq e_0 + N$. Since $p^e\alpha \geq \frac{p^{e_0}}{d} > m$, Proposition 3.4 implies that $\{p^e\alpha\} + m - 1$ is an F -jumping exponent of $\mathfrak{a}$. This gives $N + 1$ numbers in $(m - 1, m)$ that are F -jumping exponents. We deduce that there are $e_1 < e_2$ in $\{e_0, \dots, e_0 + N\}$ such that $p^{e_1}(p^{e_2-e_1} - 1)\alpha \in \mathbb{N}$, which completes the proof. $\square$

Remark 3.9. We can get a bound independent of m in the above proposition by taking $m = n$. Indeed, note first that we may assume that k is infinite: otherwise take an infinite perfect extension K of k , and replace $\mathfrak{a}$ by $\mathfrak{a} \cdot K[x_1, \dots, x_n]$ using Remark 2.18.

Now, for any maximal ideal $\mathfrak{m} \subset k[x_1, \dots, x_n] = R$, it is clear (by Proposition 2.13) that the set of jumping exponents for $\mathfrak{a}R_{\mathfrak{m}}$ is a subset of those for $\mathfrak{a}$. On the other hand, for each of the finitely many jumping exponents α in the range $[0, m]$, there is a maximal ideal $\mathfrak{m}_{\alpha}$ such that the inclusion $\tau(\mathfrak{a}^c) \subsetneq \tau(\mathfrak{a}^{c-\epsilon})$ is preserved under localization at $\mathfrak{m}_{\alpha}$. It follows from Proposition 3.4 (2) that the set of jumping exponents for $\mathfrak{a}$ is contained in (hence equal to) the union of the sets of jumping exponents for the $\mathfrak{m}_{\alpha}$, as we range over these finitely many $\mathfrak{m}_{\alpha}$.

Now, fix generators $h_1, \dots, h_m$ for $\mathfrak{a}$ of degree at most d , and let $\mathfrak{b}$ be the ideal generated by $g_i = \sum_{j=1}^m a_{i,j}h_j$ for $1 \leq i \leq n$, where $a_{i,j} \in k$. It is well-known that for every maximal ideal $\mathfrak{m}$ in R , the ideals $\mathfrak{b}R_{\mathfrak{m}}$ and $\mathfrak{a}R_{\mathfrak{m}}$ have the same integral closure if

the $a_{i,j}$ are general in k (see, for example [Laz], Example 9.6.19; note that in the proof therein one does not use the assumption that the base field is $\mathbb{C}$). Hence, by choosing the $a_{i,j}$ sufficiently general, we may assume that $\mathfrak{b}R_{\mathfrak{m}_\alpha}$ and $\mathfrak{a}R_{\mathfrak{m}_\alpha}$ have the same integral closure, and hence the same test ideals, for each of our finitely many relevant maximal ideals $\mathfrak{m}_\alpha$. It follows that the set of jumping exponents for $\mathfrak{a}$ is contained in the set of jumping exponents for the n -generated ideal $\mathfrak{b}$. Thus we get the desired statement for $\mathfrak{a}$ by applying Proposition 3.8 to $\mathfrak{b}$.

Remark 3.10. Consider several ideals $\mathfrak{a}_1, \dots, \mathfrak{a}_r$ in $k[x_1, \dots, x_n]$, where k is F -finite. The same argument used in the proof of Proposition 3.2 shows that if each $\mathfrak{a}_i$ is generated in degree at most d_i , then for every $c_1, \dots, c_r \in \mathbb{R}_+$, the mixed test ideal $\tau(\mathfrak{a}_1^{c_1} \cdots \mathfrak{a}_r^{c_r})$ can be generated in degree at most $\sum_{i=1}^r c_i d_i$. However, this assertion does not seem to have such strong consequences in the case of several ideals. The most optimistic expectation in this case is that for every $b_1, \dots, b_r$, the region

$$\{c \in \mathbb{R}_+^n \mid c_i \leq b_i \text{ for all } i\}$$

can be decomposed in a finite set of rational polytopes with non-overlapping interiors, such that on the interior of each face of such a polytope the mixed test ideal $\tau(\mathfrak{a}_1^{c_1} \cdots \mathfrak{a}_r^{c_r})$ is constant.

If k is contained in the algebraic closure of a finite field, then we can find a finite field k_0 such that every $\mathfrak{a}_i$ is generated by polynomials in $k_0[x_1, \dots, x_n]$. In this case, it follows that every $\tau(\mathfrak{a}_1^{c_1} \cdots \mathfrak{a}_r^{c_r})$ is generated by polynomials with coefficients in k_0 . In particular, if the c_i are bounded above, then the generators of the above test ideal lie in a finite-dimensional vector space over the finite field k_0 . Therefore given $\mathfrak{a}_1, \dots, \mathfrak{a}_r$, we can have only finitely many possible test ideals $\tau(\mathfrak{a}_1^{c_1} \cdots \mathfrak{a}_r^{c_r})$ if we bound the c_i .

REFERENCES

- [AMBL] J. Alvarez-Montaner, M. Blickle and G. Lyubeznik, Generators of D -modules in positive characteristic, *Math. Res. Lett.* **12** (2005), 459–473. 3, 5, 6
- [Bli] M. Blickle, The intersection homology D -module in finite characteristic, Ph.D. thesis, University of Michigan, 2001, math. AG/0110244. 11
- [BMS] M. Blickle, M. Mustařă and K. E. Smith, F -thresholds of hypersurfaces, preprint 2007. 6
- [ELSV] L. Ein, R. Lazarsfeld, K. E. Smith, and D. Varolin, Jumping coefficients of multiplier ideals, *Duke Math. J.* **123** (2004), 469–506. 1
- [HT] N. Hara and S. Takagi, On a generalization of test ideals, *Nagoya Math. J.* **175** (2004), 59–74. 2, 4, 8, 12, 13, 15
- [HH] C. Huneke and M. Hochster, Tight closure, invariant theory and the Briançon-Skoda theorem, *J. Amer. Math. Soc.* **3** (1990), 31–116. 8, 10
- [HM] N. Hara with an appendix by P. Monsky, F -pure thresholds and F -jumping exponents in dimension two, *Math. Res. Lett.* **13** (2006), 747–760. 2, 6

- [HY] N. Hara and K.-i. Yoshida, A generalization of tight closure and multiplier ideals, Trans. Amer. Math. Soc. **355** (2003), 3143–3174. [1](#), [2](#), [6](#), [8](#), [9](#), [10](#), [11](#), [13](#), [15](#)
- [Ku] E. Kunz, Characterization of regular local rings in characteristic p , Amer. J. Math. **91** (1969), 772–784. [3](#)
- [Laz] R. Lazarsfeld, Positivity in algebraic geometry II, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, Vol. **49**, Springer-Verlag, Berlin, 2004. [8](#), [13](#), [16](#), [19](#)
- [Lyu] G. Lyubeznik, F-modules: applications to local cohomology and D-modules in characteristic $p > 0$, Journal für die reine und angewandte Mathematik, **491** (1997), 65–130. [11](#)
- [Mu] M. Mustață, The multiplier ideals of a sum of ideals, Trans. Amer. Math. Soc. **354** (2002), 205–217. [17](#)
- [MTW] M. Mustață, S. Takagi and K.-i. Watanabe, F-thresholds and Bernstein-Sato polynomials, European Congress of Mathematics, 341–364, Eur. Math. Soc., Zürich, 2005. [1](#), [2](#), [3](#), [13](#)
- [Tak1] S. Takagi, Formulas for multiplier ideals on singular varieties, Amer. J. Math. **128** (2006), 1345–1362. [15](#)
- [Tak2] S. Takagi, An interpretation of multiplier ideals via tight closure, J. Algebraic Geom. **13** (2004), 393–415. [2](#)

FB MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, STANDORT ESSEN, 45117 ESSEN, GERMANY

E-mail address: `manuel.blickle@uni-essen.de`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

E-mail address: `mmustata@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

E-mail address: `kesmith@umich.edu`